# JOURNAL OF CYCLE RESEARCH

Vol. 4 No. 1

January 1955

# JOURNAL OF CYCLE RESEARCH

LEONARD W. WING, EDITOR

3875 Vorhies Road Ann Arbor, Michigan

CONTENTS, JANUARY 1955

Page

LATITUDINAL SLIPPAGE OF BOB WHITE, HAIRY WOODPECKERS, AND DOWNY WOODPECKER CYCLES, by Leonard W. Wing

3

The Journal of Cycle Research is published quarterly at 680 WEST END AVENUE. NEW YORK 25. NEW YORK. by

# THE FOUNDATION FOR THE STUDY OF CYCLES

The Foundation is a not-for-profit corporation, the board of directors of which consists of the following committee:

### UNITED STATES OF AMERICA

Copley Amory, Chairman Emeritus

Charles Greeley Ablot Research Associate Smithsonian Institution

Harold Elmer Anthony Chairman and Curator Department of Mammals American Museum of Natural History

George Backeland, Chairman of the Committee of the Foundation The Bakelite Corporation Marold Gardiner Bowen Vice Admiral, U.S. Navy, Retired Executive Director, Thomas Alva Edison Foundation

Verne Burnett Partner Verne Burnett Associate

Hon. W. Cameron Forbes
Member, Board of Trustees
Carnegie Institution of Washington;
Life Member of the Corporation,
Massachusetts Institute of Technology

Webster N. Jones Vice President Carnegie Institute of Technology

Daniel Trembly MacDougal Former Director Department of Botanical Research and Laboratory of Plant Physiology Carnegie Institution of Washington

John Quincy Stewart Professor of Astronomical Physics Princeton University

# CANADA

Hon. Charles Camsell Former Commissioner Northwest Territories Frank Cyril James Principal and Vice-Chancellor McGill University

# GREAT BRITAIN

Sir Patrick Ashley Cooper Governor Hudson's Bay Company Julian Sorell Huxley Former Director General United Nations Educational, Scientific and Cultural Organizations

Edward R. Dewey, Director

Subscription to the Journal of Cycle Research costs \$5.00 a year. Single copies are \$1.25 each. Send your check to the publication office, Foundation for the Study of Cycles, 680 West End Av.

New York 25, New York.

# LATITUDINAL SLIPPAGE

# OF BOB-WHITE, HAIRY WOODPECKER, AND DOWNY WOODPECKER CYCLES\*

BY LEONARD W. WING

RESEARCH ASSOCIATE, FOUNDATION FOR THE STUDY OF CYCLES\*\*

7 FAR-TO-YFAR changes in the numbers of animals have generated a great deal of perplexity and confusion. Field biologists studying bird and mammal life perforce have a greater interest in fluctuations, cyclic or otherwise, than almost any other group. Some hold that cycles are real, although not all changes are cyclic. Some hold, it would seem, that almost any change otherwise unexplainable must result from the workings of some mysterious thing called a "cycle." Some non-field liologists hold that there really are no such things as cycles, all fluctuations being random ones. The truth seems to be that there are cyclic changes, noncyclic (and nonrandom) changes, and random changes. I think that every natural time series probably shows all three.

Much of the confusion clearly originates in the many facets of the "cycle problem." It may be likened to the parable of the llind men and the elephant. Each student must of necessity judge from his own observation point—one might say from his own touching of the elephant.

Yet in dealing with the complexity of cycles, one must keep in mind that field biologists studying bird and mammal life recognize the fact that cycles exist. With much truth, it has been said that in this group of biologists, there is more "cycle horsepower" than in any other.

One may justly wonder why with such ample field evidence for the reality of cycles as a normal phenomenon anyone should relegate all to the dump heap of random happenings. Yet there are probably a number of reasons for this, the chief one perhaps being lack of experience with the phenomenon in the field. One studying cycles wholly on the basis of mathematical concepts quickly loses touch with reality. Hence conclusions so reached may be interesting ones but not necessarily very sound ones.

It might be well if I returned again to the question of what is or is not a cycle. Though the dictionary word cycle may mean any movement that returns to its previous state, such a broad use of the word clearly confuses more than it clarifies. With such a meaning, any fluctuation that goes up and down or down and up, in and out or out and in, and back and forth or forth and back would be a cycle. But few if any field biologists would use the term in such a loose way. A cycle as used here and

<sup>\*</sup>PART OF A PAPER PRESENTED AT THE AMERICAN ORNITHOLOGISTS' UNION MEETING, MADISON, WISCONSIN, SEPTEMBER 8 - 12, 1954.

<sup>\*\*</sup>THE INSTITUTE OF HUMAN BIOLOGY OF THE UNIVERSITY OF MICHIGAN SUPPLIED LABORATORY SPACE AND OTHER ASSISTANCE.

as used in field studies means a fluctuation that happens again and again at reasonably regular time intervals.

In order to reduce confusion, I have adopted a definition that seems exact and to reflect actual conditions in nature. A cycle may be defined as a fluctuation having (1) length, (2) rhythm, (3) timing, and (4) amplitude. Without all four, there is no cycle; the changes are simply non-cyclic fluctuations. It must be recognized, however, that at any moment we may not always have the measures for each of these. Pecause of the confusion and inexactness in use of the word cycle, I sometimes use the term rhythm, wave, or ultra-long wave for rhythmic fluctuations.

Further confusion arises in the selection of turning points used in cycle study. About the crudest form of cycle study is to pick some highs or lows in the graphed data as turning points to count in getting an average length. Equally crude is the picking of all highs and lows as turning points of cycles. While any and all are turning points of the time series, they may or may not be turning points of a cycle. In fact, they may very likely be turning points of several cycles, random events, or blended events. They may or may not be turning points of cycles; they may or may not be of random origin. They may be real for some cycles and by that token random so far as others are concerned. If one defines turning points of cycles as the high value with lower values on either side or vice versa, further progress in cycle study has been practically forbidden by definition. (If this criterion had been applied to temperature for example, we would never have discovered the cycle of the year!)

Cycles of weak amplitude by their very weakness may in a short series give the appearance of random strength and weakness. In fact, with present techniques it may be well-nigh impossible to demonstrate net rhythm in a cycle of weak amplitude. Yet gross rhythm may be clearly present.

One fact, however, does appear in the crude use of all turning points in a time series as turning points of cycles, unwittingly though it may be. If cycles are all harmonics or other mathematical relatives of each other, the manifest cycle

created by them will have mathematically related turning points. The fact that they apparently do not\* strongly contradicts the concept that all are harmonics or other "descendants" of some master length. It does not test, of course, the possibility that there may be groups of related cycles or families of cycles. It gives no test, moreover, of the reality of cycles in the time series.

Mathematical tests of cycle or rhythm significance are badly needed and no doubt will someday be devised. When they have been devised, the confusion resulting from such lack of mathematical procedures for testing or demonstrating rhythm will be removed. Statistical procedures developed for nonrhythmic situations are not now applicable as tests of rhythm and rhythm significance. Hence, one must still rely upon other procedures.

The inherent roughness of so many field data hampers cycle study. Counts of animals in the field lack a great deal of preciseness; surely this must always be so. Yet even a thermometer or rain gauge is a crude instrument. A physicist in his laboratory may measure temperature to a very close tolerance. No weather instrument is accurate to a fraction of this. Yet one soon realizes that many cycles are far too strong and prominent not to show through a considerable roughness of data.

The greatly complicated nature of cycles requires a large amount of calculation in any analytical work. It took about a quarter of a million handlings of figures, for example, to measure the cycles reported here for the three species mentioned. A previous paper dealing with global pattern of but one cycle in temperature took more. Obviously then, one limitation to measurement of cycles is the laborious computations involved. Hence, it takes a concerted, continuous, and intensive effort.

Cycle Characteristics

In addition to rhythm, timing, length,

<sup>\*</sup>COLE, LAMONT C., 1951, POPULATION CYCLES AND RANDOM OSCILLATIONS. JOURNAL OF WILD-LIFE MANAGEMENT, 15: 233-252.

RANDOM POPULATION CYCLES. I BID. 18: 2-24.

and amplitude characterizing cycles, a number of additional qualities appear. I shall mention some that bear upon the bird population cycles described in this paper.

A global behavior pattern of the 4.222-year pattern in temperature and tree rings has already been reported.\* The present paper now deals with three cycles in three bird species of the United States and Canada. The appearance of pattern on the continental and regional scope parallels that on a World basis. One may justifiably wonder if these continental expressions are not parts of global patterns.

The latitudinal phase charts of state data indicate clearly a 4.222-year rhythm that matches the global rhythm of temperature. Whether cause and effect, of course, we have no way of knowing. But a relationship, direct or indirect, is indicated. The phase chart indicates also that latitudinal slippage or passage of the epochs takes place in the bird populations as in tree rings and temperature. The rate of slippage seems to be the same in all the events reported. Though I do not know vet whether or not the 4.418-year and 3.767year lengths form global patterns in temperature. I am inclined to think that they do. We would expect therefore that timing, rhythm, and slippage shown in the three birds matches appropriate regional parts of global patterns. Some day we may know for sure.

Cycles so far measured characteristically tend to show greater amplitudes in the higher latitudes than in the lower ones. While the latitude covered by the region here studied is but about 26° and despite crude measurements of bird populations, it appears that this characteristic holds true for birds also.

# Source of Data

The lengths used in this study came originally from a study of the flight years of the Pine Grosbeak into the Lake Region and subsequent examination of global pattern in temperature. The flight years

\*WING, LEONARD W., 1954, GLOBAL PATTERN OF 4.222-YEAR CYCLES IN TEMPERATURE, JOURNAL OF CYCLE RESEARCH, 3:55-83. proved to result from a reversing cycle of 3.94-years. One member of the pair measured 4.222-years. So far as I know now, this is a single wave length or appears as a single one in a thousand years of tree rings (247 sections). The reversing mate of the 4.222-year cycle measured 3.69-years long. Further study of the 3.69-year length, however, shows that it itself is another reversing cycle. One component proves to be 3.767-years long; I have not measured its reversing mate as yet.

During the course of identifying the 4,222-year length in temperature and other events of the Northland, something about 4.4-years length conflicted with the clarity of the 4.222-year cycle. This other cycle has proved to be 4.418-years long. As long records of tree rings were used in measuring these lengths, it seems best to present the operations involved at another time as part of a separate study of rhythms in tree rings.

The data of bird abundance came from the Bird-Lore and Audubon Magazine Christmas censuses for the years 1900-1939. I converted the original figures to a ratio of birds of each species reported for each hour of censusing. This ratio of birds per hour has been used directly in the periodic tables. No moving average or other manipulation has influenced them. Cycles other than those being tested have not been minimized. As explained in other reports, J often use a modified median (Sometimes called average median or modified mean) for the respective columns of a periodic table. In the nine sections (at most) of the present periodic tables, any random distortions that may be present do not have much opportunity to offset each other. At times (though not in the tables here) I have elected to apply the principle of limited data when many apparent distortions are present. (That is, distortions so far as the cycle being tested is concerned.)

The tree ring data have been obtained from several sources as indicated in an earlier paper (Journal of Cycle Research, 3:55-83)

The quality of the data, it should be obvious, leaves much to be desired. The Christmas censuses; however, are the only abundance counts covering both a great area and a long period of years. Though the

censuses began in 1900, few regions have regular reports for the earlier years.

Several years ago, I prepared the first forty years of the censuses on the basis of birds per hour of censusing and frequency of occurrence in each state. It is this first forty years that I have used. It would be most important to use the counts since 1939, but they are available only in their untabulated hence unusable, state I see no immediate prospect that this great storehouse of data will be processed for use.

Ordinarily, I do not use a record as short as twenty to forty years in the study of cycles that are as long as three to four years. But in this instance, I have done so because the length of the cycle has been taken from a study of much longer records (up to a thousand years of tree rings). Here the data are used only to test for possible behavior patterns of the respective cycle lengths in bird numbers.

One must therefore employ the accepted scientific axiom that patterned behavior in nature betokens the operation of principles not accidents, of order not disorder, of law not chaos. The phase chart is a graphical device for showing patterns of time and place. This paper is a report on patterns of phase behavior in the Bobwhite, Hairy Woodpecker, and Downy Woodpecker.

### Bob-white

The Pob-white (Colinus virginianus) ranges over eastern North America from southern Ontario southward into Mexico. It has one characteristic that simplifies consideration of population changes; the bird is a resident species, not a migratory one. Hence, the complicating factors of seasonal movements for all practical purposes are eliminated. Few Bob-whites ever die more than a mile or so from their hatching place. The shifting of populations long has been suggested as a factor in population changes. But Bob-white numbers for analytical purposes may be considered like tree rings, crop yields, varves, or other manifestation of a nonmobile nature: whatever influenced them did so at the site. It is not a matter of the birds having moved in from some distant point.

The Christmas census data of the Bob-

white were summed for eastern North America by regions as follows (estimated general latitude in parenthesis): New England (43°N.) North Atlantic (41°N.)

Maine
New Hampshire
Vermont
Connecticut
New York
Pennsylvania
New Jersey
Maryland

Rhode Island District of Columbia
Massachusetts

South Atlantic (36°N.) Gulf (30°N.)

Virginia Georgia

West Virginia Florida

Kentucky Alabama

Tennessee Mississippi

North Carolina Louisiana

Lake Region (43°N.) Northern Plains (46°N.)

Texas

Ontario Iowa
Michigan Nebraska
Wisconsin North Dakota
Ohio South Dakota

Indiana Illinois Minnesota

South Carolina

The data for most of the states are rather meager at best. The more continuous ones were used later in a state by state phase chart. Hence, not all the thirty-four used in the region groups could be used by themselves.

As I have said before, the shortness of the series does not allow determination from internal evidence whether or not cycles of the respective lengths (3.767years, 4.222-years, and 4.418-years) are present. But a periodic table for each length for each series will indicate by strength and weakness when highs and lows would fall if present. Such show of strength or weakness is subject, of course, to distortions caused by the actions of any other cycles present, to random events, or to any other action upon the series. If the show of strength and weakness marks actual highs and lows of cycles of the respective lengths, however, their timing would be patterned.

Tables 1, 2, and 3 reproduce the periodic tables of the several region groups. They have been graphed in Figures 1, 2, and 3. The high or low of each periodic table has been measured as the midpoint between the appropriate crossing points of the curve made by the plotted means or

modified medians of the tables. As a recent base year of the 4.418-year periodic table projected forward falls at the calendar time of 1951.672, this base year added to the position of a high, as determined in Figure 1, gives the current time of ideal high. Naturally, this time of ideal high is still subject to any unneutralized influences in the table itself.

An example may make the techniques clear. The North Atlantic series gives enough data for nine sections of 4.418-years each. The mean value of the first, second, third, and fourth years after base (Table 1) are 0.37, 0.35, 0.32, and 0.39. The mean of the highest (0.39) and lowest (0.32) amplitudes is 0.355. This then lecomes the axis of the curve for the purpose of determining the midpoint of the crossings of the positive phase, which midpoint lecomes the time of high. The horizontal arrow of the North Atlantic curve marks this axis; the vertical arrow marks the midpoint at position 4.850. By adding 1951.672 to this position 4.850, the resulting 1956.522 gives a mathematical rendition of the current calendar time of ideal high. Converted to calendar days, it becomes July 10, 1956.

In a like manner, I determined the calendar times of ideal high for the Northern Plains, New England, Iake, South Atlantic, and Gulf regions for the 4.418-year cycle length. Using corresponding periodic tables, I did likewise for the 4.222-year and 3.767-year cycle lengths.

I then constructed a latitudinal phase chart for each of the cycle lengths (Figure 4), plotting the respective times of ideal high against the appropriate average latitudes. The key and current times of ideal high will be found in Table 4. To the chart for the 4.222-year cycle (middle section of Figure 4), I have added as a dashed line the temperature low of the 4.22?-year cycle from its global pattern (Journal of Cycle Research, 3:72-73, Figure 11). I have added also the epochs of the cycles in tree rings from the previous paper. In this paper the term epoch is used to indicate a time of either high or low. The tree rings form a most important corroborator of the lirds. Whichever of several different series of tree ring records are used, the epochs of corresponding latitudes fall in time with those of the

forty-year or shorter record of the Bol-white. The 4.222-year cycle has any of 50.66 months in which the epochs could fall. But all the epochs fall in a band less than sixteen months wide, which indicates both gross rhythm and a behavior pattern clearly not of random nature.

The 4.418-year cycle shows a similar behavior pattern with a spread of but thirteen of the possible fifty-three months. That of the 3.767-year cycle is also similar and has a spread of but nineteen of the forty-five possible months. The broken lines have been added to show possible slippage of the cycles, just as in the 4.222-year phase chart. Were longer records available, a narrower spread could be expected. And were it possible to neutralize or to minimize the effects of other cycles, or to eliminate local influences and randoms, the band would presumably approach the theoretical line instead of a band.

The phase charts of Figure 4 show that all three cycles have gross rhythm and pattern of timing, which suggest that the lengths are true cycles present in the Bobwhite population fluctuations. They also suggest that latitudinal slippage is present just as in the tree rings and, for the 4.222-year cycle, as in temperature.

With the knowledge that the regional groupings showed gross rhythm and patterned behavior for each of the cycles, I decided to test further with the data for each state separately. The state data for many states make but two to four sections. Though I would consider nine sections as rather inadequate at best, fewer sections are certainly worse. Nevertheless, they are all that we have available.

Yet despite this fact, the state data do show patterns in timing and gross rhythm. Tables 5, 6, and 7 are periodic tables for the 3.767, 4.222, and 4.418 year lengths, respectively, And Figures 5, 6, and 7 graph the respective averages or modified medians of the several tables. As before, the times of high (upward arrows) or low (downward arrows) are determined from the midpoints between the crossings of the axes (horizontal arrows) by the curves themselves. Adding the mathematical position of the high in the table to the most recent base year gives a representation of the calendar date of the current

time of ideal high. The current times of ideal high I have plotted on separate phase charts (Figures 8, 9, and 10). The key to the designations and times of epoch are given in Table 8.

These three phase charts show a concentration of highs in a band covering thirty-four of the possible forty-five months of the 3.767-year cycle, thirty-two of the possible fifty-one months of the 4.222-year cycle, and forty of the possible fifty-three months of the 4.417-year cycle.

But if we credit six months of the spread to the latitudinal slippage of the 3.767-year cycle, seven months to that of the 4.222-year cycle, and eight months to that of the 4.418-year cycle, the spread is reduced to twenty-eight, twenty-five, and thirty-two months of the possible forty-five, fifty-one, and fifty-three months, respectively.

If we now do likewise and credit the same amount of slippage to the same cycles for the regional Bob-white picture (Figure 4), the spread is reduced to thirteen, nine, and eleven of the possible forty-five, fifty-one, and fifty-three months, respectively. The increased concentration of regional versus state data indicates rather clearly that much of the spread from the theoretical straight line is probably induced by the shortness of record, paucity of data, and influences of local nature.

If we consider the distribution of the majority of times of ideal high, we find that they fall in but twelve of the possible forty-five months for the 3.767year cycle. Similarly the majority fall in but eight of the possible fifty-one months for the 4.222-year cycle and but seventeen of the possible fifty-three months for the 4.418-year cycle. If as before we credit six months, seven months, and eight months for latitudinal slippage of the respective cycles, the spread becomes five of a possible forty-five months for the 3.767-year cycle, one of a possible fifty-one months for the 4.222-year cycle, and nine of a possible fifty-three months for the 4.418-year cycle.

If these highs and lows occur at random, they should be equally distributed over forty-five, fifty-one, and fifty-three months. But that they are not distributed at random but are concentrated in definite parts of the respective forty-five, fifty-one, and fifty-three months is clearly powerful evidence for a rhythmic influence. It seems clear also that this spread would be reduced still further with longer records. On the basis of experience, I would look for a halving or quartering of the spread were century-long records available. Such longer records would give more opportunity for random distortions to balance each other and for other cycles to be minimized. The cycles in question would thus have greater opportunity to show through.

# Hairy and Downy Woodpeckers

Next after the Bob-white, I chose the Hairy Woodpecker (Dendrocopus villosus) and Downy Woodpecker (Dendrocopus pubescens) for a study similar to that of the state by state one for the Bob-white. But because of labor involved, only the 4.222-year cycle has been tested thus far for these two species.

I selected these two species because (1) they inhabit all states and provinces covered by the Christmas censuses, (2) they are migratory in the northern parts of their range or irregularly so, (3) they are easily observed and counted by censustakers (4) they are common enough to give substantial observational information, and (5) they are related closely enough taxonomically and ecologically to serve as comparisons with one another.

The times of ideal high have been determined by periodic tables (Tables 9 and 10), just as for the Pob-white. The means and modified medians have been graphed (Figures 11 and 12). From the times of ideal high, I constructed latitudinal phase charts separately for the Downy Woodpecker and Hairy Woodpecker (Figures 13 and 14). Table 11 gives the key to the symbols used in the phase charts and the times of epochs.

A rather different and more complicated pattern appears in these two species as compared to that of the Bob-white. The same tendency to form bands appears in the phase charts. But the highs of the southern part of the ranges cluster near the slippage line for the ideal time of low of the 4.222-year cycle in temperature as taken

from the 4.222-year temperature latitudinal phase chart (Journal of Cycle Research, 3: 72-73, Figure 11).

The band spread for the Downy Woodpecker is about eight months between latitudes 28 oN. and 38 oN. Four months of this very properly may be charged to latitudinal slippage. As in the Bob-white, if the several highs fall at random, they would have fifty-one months over which to be distributed instead of four months. Northwards from about 44 oN., the highs cluster near the line marking the slippage of the ideal time of high for the 4.222-year cycle in temperature. Because of the few areas involved, the spread cannot be determined very well, but it is greater than southward from 38 oN.

Between 38°N. and 44°N., the highs cross over from the line of temperature lows to the line of temperature highs. Just how the cross-over takes place, I cannot say. It may be that in this region, the birds show tendency for a doubling of highs. Or it may be that the state or province is too large an area for revealing the workings of this phenomenon. British Columbia and Alberta show some divergence from the pattern set by more interior areas. But the few census reports may account for it.

If one looks at it the other way around, he will note that the highs in the southern part of the range avoid the line of temperature high (if projected); the highs in the northern part similarly avoid the line of temperature lows (if projected).

Rhythm and timing are clearly present, and we may feel confident that the 4.222-year length is a true 4.222-year cycle in these birds, just as in the Bob-white. In addition, we find a difference in response to the same 4.222-year cycle that evidently is of an environmental or regional nature. It shows that birds, even of the same species and though showing the same cycle, do not necessarily show the same epochs everywhere.

The pattern in the latitudinal phase chart of the Hairy Woodpecker is identical to that of the Downy Woodpecker with a few minor variations. The crossing over takes place between about 390N. and 450N. instead

3875 Vorhies Road Ann Arbor, Michigan of 38°N, to 44°N. But this may be only an apparent difference because of paucity of data. The spread of the cluster along the line of the latitudinal slippage tends to be greater. It averages about eighteen months (fourteen net with allowance for slippage itself). Presumably the fewer numbers of Hairy Woodpeckers and consequent less adequate data than for the Downy Woodpecker plays a part in the less well defined pattern of the former species.

# Summary.

A cycle is defined as a fluctuation having (1) length, (2) rhythm, (3) timing, and (4) amplitude.

Christmas census data (as birds per hour of censusing, 1900-1939) for the Bob-white, Hairy Woodpecker, and Downy Woodpecker in the United States and Canada were tested for presence of three cycle lengths: 3.767, 4.222, and 4.418 years.

The Bob-white population data evidence presence of the three cycles in (a) regionally summed and (b) in state-summed data. Latitudinal phase charts for each cycle length show that the cycles have latitudinal slippage on a regional scale.

Though the epochs of the three cycles have any of forty-five, fifty-one, or fifty-three months, respectively, in which to fall if of random origin, all concentrate in a restricted part of these possible intervals.

The Hairy and Downy Woodpecker data from the Christmas censuses were tested for presence of the 4.222-year length.

The population highs of this cycle for both species cluster at the temperature low of the 4.222-year cycle in the southern part of the ranges and at the temperature high in the Northern part. A cross-over occurs between about Latitudes 38°-39°N. and 44°-45°N.

Regional or environmental influences may make different responses to the same cycle in the same species.

The latitudinal slippage of the Hairy and Downy Woodpeckers is the same as that found previously for the global pattern of temperature.

It is presumed that the pattern shown in the United States and Canada for the three birds may be regional expressions of appropriate global patterns.




Figure 1. Graphs of periodic table of the 3.767-year length to locate times of high in the regional data of the Bob-white. The position of the high in the periodic table added to the most recent base year gives a mathematical interpretation of the current time of ideal high. Horizontal arrows mark the axis of the curve; vertical arrows mark the point of high midway between the crossing points of the curve on its own axis.

Figure 2. Graphs of periodic tables of the 4.222-year length to locate times of high in the regional data of the Bob-white.

Note: Many of the tables and figures have been reproduced from the original work sheets.





Figure 3. Graphs of periodic tables of the 4.4]8-year length to locate times of high in the regional data of the Bob-white.



The current times of ideal high mark a well defined rhythm for each cycle length. The half-circle symbols in the 3.767-high of tree rings in British Columbia (about line of the middle figure marks the slippage of the Figure 4. Latitudinal phase charts of the regionally summed data of the Bob-white in eastern North America. 560N.) and are a part of the lower figure. The dashed the 4.222-year chart (middle figure) mark the position of 4.222-year low of temperature. Symbols are as in Table 4.




Figure 5. Graphs of periodic tables of the 3.767-year length to locate times of high of the Bob-white state data.




Figure 6. Graphs of periodic tables of the 4.222-year length to locate times of high of the Bob-white state data.




Figure 7. Graphs of periodic tables of the 4.418-year length to locate times of high of the Bob-white state data.






Figure 8. A latitudinal phase chart of the 3.767-year cycle for the Bob-white shows that the current times of ideal high fall in a band covering thirty-four of the possible forty-five months. The majority fall in a twelve month band. The dashed line marks the possible latitudinal slippage. Symbols are as listed in Table R.



eight months band. The solid slanting line marks the passage of the temperature high and the dashed slanting line the low. Symbols are as listed Table 8. ideal high fall in a band covering but thirty-two of the possible fifty-one months. A majority fall within an Figure 9. A latitudinal phase chart of the 4.222-year cycle for the Bob-white shows that the current times of



ideal high fall in a band covering forty of the possible fifty-three months of the 4.418-year cycle. The majority fall within a seventeen months band. The dashed line marks the possible latitudinal slippage. Symbols are Figure 10. A latitudinal phase chart of the 4.418-year cycle for the Bob-white shows that the current times of as listed in Table 8.




Figure 11. Graphs of periodic tables of the 4.222-year length to locate times of high of the Downy Woodpecker state data.






Figure 12. Graphs of periodic tables of the 4.222-year length to locate times of high of the Downy Woodpecker state data.



highs farther north. Retween alout 38°N. and 44°N., the Downy Woodpecker highs cross over from concentration at the temperature lows to concentration at the temperature highs. The arrows mark the high of the 4.222-year cycle high cluster near the slippage line of temperature low in the more southerly latitudes and near the temperature in British Columbia tree rings but 40 south of its actual position, which would be off the top of the chart. Figure 13. Latitudinal phase chart of the 4.222-year cycle in the Downy Woodpecker. The current times of ideal Symbols are as in Table 11.



perature lows to concentration at the temperature highs. The pattern is the same as for the Downy Woodpecker Figure 14. Latitudinal phase chart of the 4.222-year cycle in the Hairy Woodpecker. The current times of ideal high cluster near the slippage line of the temperature low in the more southerly states and the temperature high in more northerly ones. Between 390N. and 450N., the Hairy Woodpecker highs cross over from concentration at the tem-(Figure 13). The symbols are as in Table 11.

(-	TABI	LE I,	3.76 Bob.	7-year	Bur	dei To	Hou	<b>ا</b>				TAL	BL E	<i>2</i> ) ;	4, 222 3-AJJ	-year	Pen Benda	odis To perho	-+4		
7	/		Chai	stina	o Cen	anne	_								Christin	vias Ce	naus	_			
			il set			-								whole	uete	(11:00					
Bose	yes	us aj	the by	se.	6.7	bear	safte	bese				ize	ars of	ter	ose			years	agter	Base	
13000	/	2	3	3.76	/	2	3	3.76	7		Base	1	2	3	4	4,222	/	2	3	4	4,222
10	Nei	U Engla	nd .	1301	Bu	y Stal	30	N			1900	42	Peury	Carro S	450 N	1		103	on and	uter	410 N
1901	.09	.02	102	.06	0	290	2.55	2.00	1		1904	.05	.07	.02	02	.06	133	.04	.05	120	1/2
1909	187	.17	103	02	,51_ ©	.31_	.34	(25)			1913	.32	.08	,21"	1/2		143	131	.69	.97	
1916	112	10	.08	107	+37	.0	.09	3 .0			1921	.10	.09	13	107	122	159	.50	.29	138	.26
1927	.07	,22	116	99,	015	137	177	050			1930	.04	.01	20.	.06		133	.69	.40	172	
1927	.01	.03	.06	.04	.47	126	-84	29			1938	1.12	1.00	168	. 74	-	325	3.47	2.49	3.90	
ave	87	1.07	.93	.80	4.87	6.56	5.46	7.98			ave	./3	.//	.07	.08		.39	، عرب ا	127	143	
1001	Nast	n ottle	ente	410 N	مّه	10 Stal					1900	Sau	et all	unter	36°N		9	web St	etes	30°/	r.
1901	104	115	.20	15	. 70	.32	1.38	.62			1904	1,24	1.85.	170	,93	1.77		35	,34	- ي د.	
1909	33	.52	.31	.69	149	1.17	.68	1122			1917	.23.	.87	1.25	1.00		.5-3	-02	.09	.57	
1916 1920 1914	.51	143	173	129	1.66	1.38	. 84	1.45			1926	1.980	.83	1.860	2.60 -	.66	145	175	.47	.70	.31
1917	200	.28	122	150	1.66	1.24	.56	150			1930	-36	.82	1.860	1.07		126	147	126	.481 .7h	
1951 1955	.16	126	.71	137	.46	.60	1.42	1.34			1938	£ 5.80	3.82	5119	6.42		3.15	2.20	200	3.08	_
aure	3.90	3.72	3.88	3.72	6.70		7.94	8.48			ave.	.47	164	187	1:08		052	.37	-33	151	
1901	Sau	1. O	Clarte	369	W 0-		h0 .				1900		-71	,20	.34		Ŋ	archi	, pla	ins 4	LON
1965	1.24		.70	(93)	1/21	then 1	Planis	46°N.			1909	1.17	166	168	1138	190				: 1	
1909	1.00	1230		185							1913	34	164	1.22	.48		~	122	.03	104-	
1916 1920 1924	192	1.56	1.98	194	-01	.54	,44-	150			1921	1.38	184	1.45	1.66	192	154	131	.43	:76	.97
1937	.83	1.07	1.87	.36.	-31	143	.76	126			1930		.86	.83	1.45		116	:3/	,26	174	
1937	640	.41	1.07	.42	-31	183	.74	.69			1938	7.51	5.95	6,88	7,20		2.18	1.40	2.61	2.14	
ave	.92	1.07	5.67	5.23	.92	3.03	2.94	192				, 8 3	166	.76	.80		.44	-28	141	.43	
	1		U	1110 (4	່.	1-7	7.4	1 1				ì			1	Ì.		l i	]		
(	Table		Bock	6-Whi restrone	of Ben to Bu	odi 1	Toble have				Rest							ROUPS OF			gyola
			Bo	6-Who	to , Bu	oder 1.	Totle				Morth	on Group arn Flai	Latitu	de Symbo	01 3.767-	-year oy	10 4.22	2-year o	yole 4.4	18-year	cycle
Bu			Bo	6-Whi restrone	to , Bu	oder 1	Jede hase				North New E: Laks	on Group ern Plai ngland	Latitu ns 46 43		01 3.767-	-year oy	10 4.22	2-year o	yole 4.4	18-year	ayale 1955 1955
	W Y	ado a	Bo Ch	6-Whi without 0 ml Bese 4	to , Bu	odie 1	year	o agte	ti Stab	4.418	North New E: Lake North	on Group arn Flai	Latitu ns 46 43 42 c 41	de Symbo	01 3.767-	-year oy	10 4.22	2-year o	yole 4.4	18-year	oyole 1955 1955 1956 1956 1955
1899	W Y	ars o	Che Charles Sand S	b-Whose whose Besse y Lates	4.418	ido-pa	year 2	a ple	tistat		North New E Lake North South Gulf	on Group ern Plai ngland Atlanti Atlanti	Latitu ons 45 43 42 8 41 9 36 30	de Symbolica de Sy	ol 3.767- Oct. Feb. Feb. Feb. Dec.	7, 1955 3, 1952 1, 1953 1, 1952 12, 1954 19, 195	Jar Apr Apr Mar Sey 2 Mar 3 Jur		1701e 4.4 155 De 1956 Oc 155 Ar 1955 Je 1955 Be		ayale 1955 1955 1956 1956 1955
1899 1903 1907 1912	W 44	2 2 2 2 2 2 2 2 32.0	Book of the stand	Besse 4	4.418	ido-pa	year	0 apte	#15tab	,05	Morth New E Lake North South Gulf Tree Briti Vermon	on Group ern Plai ngland Atlanti Atlanti ring (S	Intitu ons 46 43 42 c 41 o 36 30 ubject t	de Symbolica de Sy	ol 3.767- Oct. Feb. Feb. Feb. Des.	7, 1956 3, 1952 1, 1953 1, 1952 12, 196 19, 195	Jar Apr Mar Ser Mar Jur Ser Feed	22-year of a. 28, 19 11 24, 1 10h 6, 19 0t. 12, 1 10h 13, 1 10e 12, 19 0noiliati	1901e 4.4 1955 De 1956 Oc 1955 Ar 1955 Be 1955 Oc 1001)	18-year 6. 13, 1 10. 16, 1 11, 10, 1 11, 10, 1 11, 10, 1 11, 10, 1	1955 1955 1956 1956 1956 1955
1899 1903 1907 1912	/ / / / / / / / / / / / / / / / / / /	2 Em Em . 06 . 132 . 09	gland S land S land S	bese Sese 101 107 107	4.4/8	ido-pa	year  Ant  11  12  12  13  12  15  10  10  10  10  10  10  10  10  10	0 apte 103 104 109 131 131 139	105 .05 .05 .05 .05 .05 .05 .05 .05 .05 .	.05	Morth New E Lake North South Gulf Tree Briti Vermon	on Group arn Plai ngland Atlanti Atlanti ring (S sh Column nt ylvania	Iatitu ons 45 43 42 6 41 6 36 30 ubject t bia 86 44 42 36	de Symbol 2 3 4 5 6 o growth	ol 3.767- Oct. Feb. Feb. Feb. Dec. h and cal- on Jan. Sapril Oct. *Jan.	7, 1956 3, 1952 1, 1953 1, 1952 12, 196 19, 195	Jar Apr Mar Ser Mar Jur Ser Feed	22-year on 28, 19 11 24, 1 10 6, 19 11 12, 1 10 13, 1 10 12, 19	1901e 4.4 1955 De 1956 Oc 1955 Ar 1955 Be 1955 Oc 1001)	18-year	1955 1955 1956 1956 1956 1955
1899 1903 1907 1912 1916 1921 1921	02 .02 .02 .00 .100 .101 .1Y	2 En 6 : 32	3 land S .09 .09 .09 .09 .09 .09 .09 .09 .09 .09	bese y takes 102 107 107 107 107	4.418	109 - 120 120 134 197-151 126	4 have 2 have 2 have 2 have 2 have 32	03 03 03 137 129 129 126	\$ 55.05 .05 .05 .69 .38 .58	,05	Morth New E Lake North South Gulf Tree Briti Vermon Penns	on Group srn Plai ngland Atlanti Atlanti ring (S sh Column nt ylvania	Intitu ons 46 43 42 41 6 36 30 ubject t	de Symbol 2 3 4 5 6 o growth	ol 3.767- Oct. Feb. Feb. Feb. Dec. h and cal- on Jan. Sapril Oct. *Jan.	7, 1955 3, 1952 1, 1953 1, 1953 1, 1953 12, 195 19, 195 lendar y 31, 195 13, 195 10, 195	Jar Apr Mar Ser Mar Jur Ser Feed	22-year of a. 28, 19 11 24, 1 10h 6, 19 0t. 12, 1 10h 13, 1 10e 12, 19 0noiliati	1901e 4.4 1955 De 1956 Oc 1955 Ar 1955 Be 1955 Oc 1001)	18-year 6. 13, 1 10. 16, 1 11, 10, 1 11, 10, 1 11, 10, 1 11, 10, 1	1955 1955 1956 1956 1956 1955
/899 /903 /903 /907 /916 /92/ /93/ /93/	02 .02 .02 .02 .01 .01 .04 .04 .04 .04	2 2 2 2 2 2 2 2 3 2 0 3 2 0 3 1 0 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	Bo Ch	6- Who we have a series of 107 .07 .07 .07 .07 .06	4.418 103 12 16	100-pa	4 A A A A A A A A A A A A A A A A A A A	0 2 de 10 10 10 10 10 10 10 10 10 10 10 10 10	\$ 155.69 .05 .69 .08 .58 .72 .72	.05	Morth New E Lake North South Gulf Tree Briti Vermon Penns	on Group srn Plai ngland Atlanti Atlanti ring (S sh Column nt ylvania	Iatitu ons 45 43 42 6 41 6 36 30 ubject t bia 86 44 42 36	de Symbol 2 3 4 5 6 o growth	ol 3.767- Oct. Feb. Feb. Feb. Dec. h and cal- on Jan. Sapril Oct. *Jan.	7, 1955 3, 1952 1, 1953 1, 1953 1, 1953 12, 195 19, 195 lendar y 31, 195 13, 195 10, 195	Jar Apr Mar Ser Mar Jur Ser Feed	22-year of a. 28, 19 11 24, 1 10h 6, 19 0t. 12, 1 10h 13, 1 10e 12, 19 0noiliati	1901e 4.4 1955 De 1956 Oc 1955 Ar 1955 Be 1955 Oc 1001)	18-year 6. 13, 1 10. 16, 1 11, 10, 1 11, 10, 1 11, 10, 1 11, 10, 1	1955 1955 1956 1956 1956 1955
1899 1903 1907 1912 1921 1921 1939 1938	00 10 10 10 10 10 10 10 10 10 10 10 10 1	2 Em Em (0) (0) (0) (0) (0) (0) (0) (0) (0) (0)	3 land S .09 .09 .09 .09 .09 .09 .09 .09 .09 .09	Bese 4 107 .07 .07 .07 .06 .68	4.418 103 12 16	109 - 120 109 - 120 134 197. 126 192.	4 have 2 have 2 have 2 have 2 have 32	03 03 03 137 129 129 126	\$ 55.05 .05 .05 .69 .38 .58	.05	Morth New E Lake North South Gulf Tree Briti Vermon Penns	on Group srn Plai ngland Atlanti Atlanti ring (S sh Column nt ylvania	Iatitu ons 45 43 42 6 41 6 36 30 ubject t bia 86 44 42 36	de Symbol 2 3 4 5 6 o growth	ol 3.767- Oct. Feb. Feb. Feb. Dec. h and cal- on Jan. Sapril Oct. *Jan.	7, 1955 3, 1952 1, 1953 1, 1953 1, 1953 12, 195 19, 195 lendar y 31, 195 13, 195 10, 195	Jar Apr Mar Ser Mar Jur Ser Feed	22-year of a. 28, 19 11 24, 1 10h 6, 19 0t. 12, 1 10h 13, 1 10e 12, 19 0noiliati	1901e 4.4 1955 De 1956 Oc 1955 Ar 1955 Be 1955 Oc 1001)	18-year 6. 13, 1 10. 16, 1 11, 10, 1 11, 10, 1 11, 10, 1 11, 10, 1	1955 1955 1956 1956 1956 1955
1899 1903 1907 1912 1916 1927 1930 1934 1938 1939	00 10 10 10 10 10 10 10 10 10 10 10 10 1	2 Em Em	Bock  of land  sland  of los	6- Whose of the service of the servi	4.418 103 12 16	100-pa	4 Azie 2	2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	\$ 105 105 105 151 169 138 158 172 177 3.54	.05 .5/ .22	Morth New E Lake North South Gulf Tree Briti Vermon Penns	on Group srn Plai ngland Atlanti Atlanti ring (S sh Column nt ylvania	Iatitu ons 45 43 42 6 41 6 36 30 ubject t bia 86 44 42 36	de Symbol 2 3 4 5 6 o growth	ol 3.767- Oct. Feb. Feb. Feb. Dec. h and cal- on Jan. Sapril Oct. *Jan.	7, 1955 3, 1952 1, 1953 1, 1953 1, 1953 12, 195 19, 195 lendar y 31, 195 13, 195 10, 195	Jar Apr Mar Ser Mar Jur Ser Feed	22-year of a. 28, 19 11 24, 1 10h 6, 19 0t. 12, 1 10h 13, 1 10e 12, 19 0noiliati	1901e 4.4 1955 De 1956 Oc 1955 Ar 1955 Be 1955 Oc 1001)	18-year 6. 13, 1 10. 16, 1 11, 10, 1 11, 10, 1 11, 10, 1 11, 10, 1	1955 1955 1956 1956 1956 1955
/899 /903 /903 /903 /92/ /92/ /93/ /93/ EDE, H	02 .02 .02 .09 .09 .178 .07 .09 .07	2 Em Em	Jan 3 sand 3 sand 3 son	6- Whose of the service of the servi	4.4/8 103 12 16 apris al.	100-pa	year 22 15 15 15 15 15 15 15 15 15 15 15 15 15	2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	105 105 151 169 138 158 177 147 3.59	.05 .5/ .22 apri al	Morth New E Lake North South Gulf Tree Briti Vermon Penns	on Group srn Plai ngland Atlanti Atlanti ring (S sh Column nt ylvania	Iatitu ons 45 43 42 6 41 6 36 30 ubject t bia 86 44 42 36	de Symbol 2 3 4 5 6 o growth	ol 3.767- Oct. Feb. Feb. Feb. Dec. h and cal- on Jan. Sapril Oct. *Jan.	7, 1955 3, 1952 1, 1953 1, 1953 1, 1953 12, 195 19, 195 lendar y 31, 195 13, 195 10, 195	Jar Apr Mar Ser Mar Jur Ser Feed	22-year of a. 28, 19 11 24, 1 10h 6, 19 0t. 12, 1 10h 13, 1 10e 12, 19 0noiliati	1901e 4.4 1955 De 1956 Oc 1955 Ar 1955 Be 1955 Oc 1001)	18-year 6. 13, 1 10. 16, 1 11, 10, 1 11, 10, 1 11, 10, 1 11, 10, 1	1955 1955 1956 1956 1956 1955
/899 /903 /903 /903 /92/ /92/ /93/ /93/ EDE, H	102 00 110 111 110 110 110 110 110 110 1	20 Em Em 106 : 32	100 Ch	6-25 million 200 m	4.418 103 1/2 1/6 2/42 200 270	.09 .20 .37 .5/ .33 .34 .87	" have "	203 203 203 203 203 203 203 203 203 203	\$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$	.05 .5/ .22 apriak .35½	Morth New E Lake North South Gulf Tree Briti Vermon Penns	on Group srn Plai ngland Atlanti Atlanti ring (S sh Column nt ylvania	Iatitu ons 45 43 42 6 41 6 36 30 ubject t bia 86 44 42 36	de Symbol 2 3 4 5 6 o growth	ol 3.767- Oct. Feb. Feb. Feb. Dec. h and cal- on Jan. Sapril Oct. *Jan.	7, 1955 3, 1952 1, 1953 1, 1953 1, 1953 12, 195 19, 195 lendar y 31, 195 13, 195 10, 195	Jar Apr Mar Ser Mar Jur Ser Feed	22-year of a. 28, 19 11 24, 1 10h 6, 19 0t. 12, 1 10h 13, 1 10e 12, 19 0noiliati	1901e 4.4 1955 De 1956 Oc 1955 Ar 1955 Be 1955 Oc 1001)	18-year 6. 13, 1 10. 16, 1 11, 10, 1 11, 10, 1 11, 10, 1 11, 10, 1	1955 1955 1956 1956 1956 1955
/899 /903 /903 /903 /92/ /92/ /93/ /93/ EDE, H	102 00 110 111 110 110 110 110 110 110 1	2 2 Em Em	100 Ch	6- 25 mm Base 2	4.4/8  103 11 16  aprigate	.09 .20 .37 .5/ .33 .34 .87	12 1/2 1/2 1/2 1/2 1/2 1/2 1/2 1/2 1/2 1	2 400 00 00 00 00 00 00 00 00 00 00 00 00	\$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$	.05 .5/ .22 apri al .354	Morth New E Lake North South Gulf Tree Briti Vermon Penns	on Group srn Plai ngland Atlanti Atlanti ring (S sh Column nt ylvania	Iatitu ons 45 43 42 6 41 6 36 30 ubject t bia 86 44 42 36	de Symbol 2 3 4 5 6 o growth	ol 3.767- Oct. Feb. Feb. Feb. Dec. h and cal- on Jan. Sapril Oct. *Jan.	7, 1955 3, 1952 1, 1953 1, 1953 1, 1953 12, 195 19, 195 lendar y 31, 195 13, 195 10, 195	Jar Apr Mar Ser Mar Jur Ser Feed	22-year of a. 28, 19 11 24, 1 10h 6, 19 0t. 12, 1 10h 13, 1 10e 12, 19 0noiliati	1901e 4.4 1955 De 1956 Oc 1955 Ar 1955 Be 1955 Oc 1001)	18-year 6. 13, 1 10. 16, 1 11, 10, 1 11, 10, 1 11, 10, 1 11, 10, 1	1955 1955 1956 1956 1956 1955
/899 /903 /907 /912 /916 /93/ /938 Eac. /899 /943 /905 /906 /906 /906 /906 /906	102 00 110 111 110 110 110 110 110 110 1	2 2 Em Em	100 Ch	6- Whomas of the state of the s	4.418 103 12 116 axis at 12 170 192 1.87	.09 120 139 197 120 134 133 134 137 137 137 137 137 137 137 136 137 137 137 137 137 137 137 137 137 137	12 1/2 1/2 1/2 1/2 1/2 1/2 1/2 1/2 1/2 1	2.57 2.57 2.57 2.57 2.57 2.57 2.57 2.57 2.57 2.57 2.57 2.57	\$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$	.05 .5/ .22 apriak .35½	Morth New E Lake North South Gulf Tree Briti Vermon Penns	on Group srn Plai ngland Atlanti Atlanti ring (S sh Column nt ylvania	Iatitu ons 45 43 42 6 41 6 36 30 ubject t bia 86 44 42 36	de Symbol 2 3 4 5 6 o growth	ol 3.767- Oct. Feb. Feb. Feb. Dec. h and cal- on Jan. Sapril Oct. *Jan.	7, 1955 3, 1952 1, 1953 1, 1953 1, 1953 12, 195 19, 195 lendar y 31, 195 13, 195 10, 195	Jar Apr Mar Ser Mar Jur Ser Feed	22-year of a. 28, 19 a. 28, 19 a. 24, 1 ch 6, 19 ot. 12, 1 ch 13, 1 ce 12, 19 onciliati	1901e 4.4 1955 De 1956 Oc 1955 Ar 1955 Be 1955 Oc 1001)	18-year 6. 13, 1 10. 16, 1 11, 10, 1 11, 10, 1 11, 10, 1 11, 10, 1	1955 1955 1956 1956 1956 1955
/899 /903 /907 /912 /916 /93/ /938 Eac. /899 /943 /905 /906 /906 /906 /906 /906	02 00 100 121 1.04 1.05 1.06 1.06 1.06 1.06 1.06 1.06 1.06 1.06	2 Ew Em Em (32 ° ° ° ° ° ° ° ° ° ° ° ° ° ° ° ° ° ° °	Che	6- 25 102 102 102 102 102 102 102 102 102 102	4.418 103 112 116 270 192 1.87	.09 120 139 197 120 134 133 134 137 137 137 137 137 137 137 136 137 137 137 137 137 137 137 137 137 137	Gran 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	240 233 29 29 250 246 251 250 250 250 250 250 250 250 250 250 250	155 16 105 157 157 157 157 157 157 157 157 157 15	.05 .5/ .22 agazak .35/2	Morth New E Lake North South Gulf Tree Briti Vermon Penns	on Group srn Plai ngland Atlanti Atlanti ring (S sh Column nt ylvania	Iatitu ons 45 43 42 6 41 6 36 30 ubject t bia 86 44 42 36	de Symbol 2 3 4 5 6 o growth	ol 3.767- Oct. Feb. Feb. Feb. Dec. h and cal- on Jan. Sapril Oct. *Jan.	7, 1955 3, 1952 1, 1953 1, 1953 1, 1953 12, 195 19, 195 lendar y 31, 195 13, 195 10, 195	Jar Apr Mar Ser Mar Jur Ser Feed	22-year of a. 28, 19 a. 28, 19 a. 24, 1 ch 6, 19 ot. 12, 1 ch 13, 1 ce 12, 19 onciliati	1901e 4.4 1955 De 1956 Oc 1955 Ar 1955 Be 1955 Oc 1001)	18-year 6. 13, 1 10. 16, 1 11, 10, 1 11, 10, 1 11, 10, 1 11, 10, 1	1955 1955 1956 1956 1956 1955
/899 /903 /903 /903 /92/ /92/ /93/ /93/ EDE, H	102 00 110 111 110 110 110 110 110 110 1	2 2 2 2 2 2 32 32 32 32 32 32	Che Chino China Chino China Chino Ch	6- Who without a wind of the control	4.418 103 12 116 axis at 12 170 192 1.87	.09 .09 .20 .34 .37 .57 .33 .37 .37 .37 .37 .37 .37 .37 .37 .3	12 1/2 1/2 1/2 1/2 1/2 1/2 1/2 1/2 1/2 1	2.89 2.89 2.89 2.89 2.89 2.89 2.89 2.89	\$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$	.05 .5/ .22 apri al .354	Morth New E Lake North South Gulf Tree Briti Vermon Penns	on Group srn Plai ngland Atlanti Atlanti ring (S sh Column nt ylvania	Iatitu ons 45 43 42 6 41 6 36 30 ubject t bia 86 44 42 36	de Symbol 2 3 4 5 6 o growth	ol 3.767- Oct. Feb. Feb. Feb. Dec. h and cal- on Jan. Sapril Oct. *Jan.	7, 1955 3, 1952 1, 1953 1, 1953 1, 1953 12, 195 19, 195 lendar y 31, 195 13, 195 10, 195	Jar Apr Mar Ser Mar Jur Ser Feed	22-year of a. 28, 19 a. 28, 19 a. 24, 1 ch 6, 19 ot. 12, 1 ch 13, 1 ce 12, 19 onciliati	1901e 4.4 1955 De 1956 Oc 1955 Ar 1955 Be 1955 Oc 1001)	18-year 6. 13, 1 10. 16, 1 11, 10, 1 11, 10, 1 11, 10, 1 11, 10, 1	1955 1955 1956 1956 1956 1955
/899 /903 1907 1916 1924 1930 1930 1938 Ear, 111 1930 1997 1993 1993 1993 1994 1994 1994 1994 1995 19	02 00 100 121 1.04 1.05 1.06 1.06 1.06 1.06 1.06 1.06 1.06 1.06	2 Euro E Euro E E E E E E E E E E E E E E E E E E E	Che Chine Che Che Che Che Che Che Che Che Che Ch	6- Who without the control of the co	4.418 103 112 116 270 192 1.87 4.87	.09 .09 .20 .34 .37 .57 .33 .37 .37 .37 .37 .37 .37 .37 .37 .3	June 318 3.18 3.18 3.18 3.18 3.18 3.18 3.18	2.87 3.17 7.26 1.29 1.20 2.87 1.31 3.1 7.20 1	38 .572 .69 .69 .572 .727 .89 .89 .270 .99 .89 .270 .99 .89 .270 .99 .89	.05 .5/ .22 agazak .35/2	Morth New E Lake North South Gulf Tree Briti Vermon Penns	on Group srn Plai ngland Atlanti Atlanti ring (S sh Column nt ylvania	Iatitu ons 45 43 42 6 41 6 36 30 ubject t bia 86 44 42 36	de Symbol 2 3 4 5 6 o growth	ol 3.767- Oct. Feb. Feb. Feb. Dec. h and cal- on Jan. Sapril Oct. *Jan.	7, 1955 3, 1952 1, 1953 1, 1953 1, 1953 12, 195 19, 195 lendar y 31, 195 13, 195 10, 195	Jar Apr Mar Ser Mar Jur Ser Feed	22-year of a. 28, 19 a. 28, 19 a. 24, 1 ch 6, 19 ot. 12, 1 ch 13, 1 ce 12, 19 onciliati	1901e 4.4 1955 De 1956 Oc 1955 Ar 1955 Be 1955 Oc 1001)	18-year 6. 13, 1 10. 16, 1 11, 10, 1 11, 10, 1 11, 10, 1 11, 10, 1	1955 1955 1956 1956 1956 1955
/899 /903 1907 1916 1924 1930 1930 1938 Ear, 111 1930 1997 1993 1993 1993 1994 1994 1994 1994 1995 19	00 1/0 1/0 1/0 1/0 1/0 1/0 1/0 1/0 1/0 1	2 Ew Em E 6 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	Che Chino Che Chino Che Chino Che	5- Who without the condition of the cond	4.418 103 112 116 270 192 1.87 4.87	.09 .09 .20 .34 .37 .57 .33 .37 .37 .37 .37 .37 .37 .37 .37 .3	June 318 3.18 3.18 3.18 3.18 3.18 3.18 3.18	240 233 29 29 250 246 251 250 250 250 250 250 250 250 250 250 250	38 .572 .69 .69 .572 .727 .89 .89 .270 .99 .89 .270 .99 .89 .270 .99 .89	.05 .5/ .22 agazak .35/2	Morth New E Lake North South Gulf Tree Briti Vermon Penns	on Group srn Plai ngland Atlanti Atlanti ring (S sh Column nt ylvania	Iatitu ons 45 43 42 6 41 6 36 30 ubject t bia 86 44 42 36	de Symbol 2 3 4 5 6 o growth	ol 3.767- Oct. Feb. Feb. Feb. Dec. h and cal- on Jan. Sapril Oct. *Jan.	7, 1955 3, 1952 1, 1953 1, 1953 1, 1953 12, 195 19, 195 lendar y 31, 195 13, 195 10, 195	Jar Apr Mar Ser Mar Jur Ser Feed	22-year of a. 28, 19 a. 28, 19 a. 24, 1 ch 6, 19 ot. 12, 1 ch 13, 1 ce 12, 19 onciliati	1901e 4.4 1955 De 1956 Oc 1955 Ar 1955 Be 1955 Oc 1001)	18-year 6. 13, 1 10. 16, 1 11, 10, 1 11, 10, 1 11, 10, 1 11, 10, 1	1955 1955 1956 1956 1956 1955
/899 /903 1907 1916 1924 1930 1930 1938 Ear, 111 1930 1997 1993 1993 1993 1994 1994 1994 1994 1995 19	02 .01 .02 .01 .02 .03 .07 .07 .07 .07 .07 .07 .07 .07 .07 .07	2 2 Ew Ewit 500 101 101 101 101 101 101 101 101 101	Che Chino Che Chino Che Chino Che	6- 24 Cates of Cates	4.418 103 12 116 april ad 12 198 199 191 191 191 191 191 191 191 191	.09 .09 .20 .34 .37 .57 .33 .37 .37 .37 .37 .37 .37 .37 .37 .3	12 15 15 15 15 15 15 15 15 15 15 15 15 15	202 202 202 202 202 202 202 202 202 202	15 15 15 15 15 15 15 15 15 15 15 15 15 1	.05 .5/ .22 agazak .35/2	Morth New E Lake North South Gulf Tree Briti Vermon Penns	on Group srn Plai ngland Atlanti Atlanti ring (S sh Column nt ylvania	Iatitu ons 45 43 42 6 41 6 36 30 ubject t bia 86 44 42 36	de Symbol 2 3 4 5 6 o growth	ol 3.767- Oct. Feb. Feb. Feb. Dec. h and cal- on Jan. Sapril Oct. *Jan.	7, 1955 3, 1952 1, 1953 1, 1953 1, 1953 12, 195 19, 195 lendar y 31, 195 13, 195 10, 195	Jar Apr Mar Ser Mar Jur Ser Feed	22-year of a. 28, 19 a. 28, 19 a. 24, 1 ch 6, 19 ot. 12, 1 ch 13, 1 ce 12, 19 onciliati	1901e 4.4 1955 De 1956 Oc 1955 Ar 1955 Be 1955 Oc 1001)	18-year 6. 13, 1 10. 16, 1 11, 10, 1 11, 10, 1 11, 10, 1 11, 10, 1	1955 1955 1956 1956 1956 1955
/899 /903 1907 1916 1924 1930 1930 1938 Ear, 111 1930 1997 1993 1997 1993 1997 1993 1997 1993 1997 1993 1993 1994 1994 1994 1994 1994 1994 1994 1994 1994 1994 1995 19	02 .01 .02 .01 .02 .03 .07 .07 .07 .07 .07 .07 .07 .07 .07 .07	2 2 Ew Ewit Ewit 6 132 - 09 105 101 177 198 164 177 198 164 164 179 198 164 179 198 164 179 198 164 179 198 164 179 198 164 179 198 164 179 198 164 179 198 164 179 179 179 179 179 179 179 179 179 179	Che Chino China Chino China Chino Ch	5- 24 Care 1 10 1 10 1 10 1 10 1 10 1 10 1 10 1	1,418 103 12 116 202 116 170 192 1.87 481 188	100 - pase	12 15 15 15 15 15 15 15 15 15 15 15 15 15	24 25 25 25 25 25 25 25 25 25 25 25 25 25	15 15 15 15 15 15 15 15 15 15 15 15 15 1	.05 .5/ .22 agazak .35/2	Morth New E Lake North South Gulf Tree Briti Vermon Penns	on Group srn Plai ngland Atlanti Atlanti ring (S sh Column nt ylvania	Iatitu ons 45 43 42 6 41 6 36 30 ubject t bia 86 44 42 36	de Symbol 2 3 4 5 6 o growth	ol 3.767- Oct. Feb. Feb. Feb. Dec. h and cal- on Jan. Sapril Oct. *Jan.	7, 1955 3, 1952 1, 1953 1, 1953 1, 1953 12, 195 19, 195 lendar y 31, 195 13, 195 10, 195	Jar Apr Mar Ser Mar Jur Ser Feed	22-year of a. 28, 19 a. 28, 19 a. 24, 1 ch 6, 19 ot. 12, 1 ch 13, 1 ce 12, 19 onciliati	1901e 4.4 1955 De 1956 Oc 1955 Ar 1955 Be 1955 Oc 1001)	18-year 6. 13, 1 10. 16, 1 11, 10, 1 11, 10, 1 11, 10, 1 11, 10, 1	1955 1955 1956 1956 1956 1955
/899 /903 1907 1916 1924 1930 1930 1938 Ear, 111 1930 1997 1993 1997 1993 1997 1993 1997 1993 1997 1993 1993 1994 1994 1994 1994 1994 1994 1994 1994 1994 1994 1995 19	02 .01 .02 .01 .02 .03 .07 .07 .07 .07 .07 .07 .07 .07 .07 .07	2 2 Ew Ewit Ewit 6 132 - 09 105 101 177 198 164 177 198 164 164 179 198 164 179 198 164 179 198 164 179 198 164 179 198 164 179 198 164 179 198 164 179 198 164 179 179 179 179 179 179 179 179 179 179	Che Chino Che Chino Che Chino Che	6- 24 Cates of Cates	103 12 16 april ad 102 192 1.87 april ad 1986 1.48	100 - pase	Gran 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	202 202 202 202 202 202 202 202 202 202	15 15 15 15 15 15 15 15 15 15 15 15 15 1	.05 .5/ .22 .34 .34 .47 .47	Morth New E Lake North South Gulf Tree Briti Vermon Penns	on Group srn Plai ngland Atlanti Atlanti ring (S sh Column nt ylvania	Iatitu ons 45 43 42 6 41 6 36 30 ubject t bia 86 44 42 36	de Symbol 2 3 4 5 6 o growth	ol 3.767- Oct. Feb. Feb. Feb. Dec. h and cal- on Jan. Sapril Oct. *Jan.	7, 1955 3, 1952 1, 1953 1, 1953 1, 1953 12, 195 19, 195 lendar y 31, 195 13, 195 10, 195	Jar Apr Mar Ser Mar Jur Ser Feed	22-year of a. 28, 19 a. 28, 19 a. 24, 1 ch 6, 19 ot. 12, 1 ch 13, 1 ce 12, 19 onciliati	1901e 4.4 1955 De 1956 Oc 1955 Ar 1955 Be 1955 Oc 1001)	18-year 6. 13, 1 10. 16, 1 11, 10, 1 11, 10, 1 11, 10, 1 11, 10, 1	1955 1955 1956 1956 1956 1955
/899 /903 /907 /912 /916 /93/ /938 Eac. /899 /943 /905 /906 /906 /906 /906 /906	00 1/0 1/0 1/0 1/0 1/0 1/0 1/0 1/0 1/0 1	2 2 Ew Ewit 500 101 101 101 101 101 101 101 101 101	Che Chino China Chino China Chino Ch	6- 10 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	103 12 16 april ad 102 192 1.87 april ad 1986 1.48	100 - pase	12 15 15 15 15 15 15 15 15 15 15 15 15 15	24 25 25 25 25 25 25 25 25 25 25 25 25 25	38 .572 .69 .69 .572 .727 .89 .89 .270 .99 .89 .270 .99 .89 .270 .99 .89	.05 .5/ .22 .42 .35/ .34 .34 .37 .47	Morth New E Lake North South Gulf Tree Briti Vermon Penns	on Group srn Plai ngland Atlanti Atlanti ring (S sh Column nt ylvania	Iatitu ons 45 43 42 6 41 6 36 30 ubject t bia 86 44 42 36	de Symbol 2 3 4 5 6 o growth	ol 3.767- Oct. Feb. Feb. Feb. Dec. h and cal- on Jan. Sapril Oct. *Jan.	7, 1955 3, 1952 1, 1953 1, 1953 1, 1953 12, 195 19, 195 lendar y 31, 195 13, 195 10, 195	Jar Apr Mar Ser Mar Jur Ser Feed	22-year of a. 28, 19 a. 28, 19 a. 24, 1 ch 6, 19 ot. 12, 1 ch 13, 1 ce 12, 19 onciliati	1901e 4.4 1955 De 1956 Oc 1955 Ar 1955 Be 1955 Oc 1001)	18-year 6. 13, 1 10. 16, 1 11, 10, 1 11, 10, 1 11, 10, 1 11, 10, 1	1955 1955 1956 1956 1956 1955

					-	1	AK	3 1	E	5	1										
					3.72	V_ 48	a	Per	rod	tei	toble	2 . /	Bro	f-4	hil	<u> </u>					
,	,			(	The	istr	TIL	06	ense	ese	4	Bu	rdo,	eer 1	rous	-					
	- ,				1										<u> </u>				1		
Bese	y	rars	yes	base	(20)	l u	Dalo	Mest	are (	420)	Bese	: ge	ars. a	tu.	Lase		ge	as	after	base	
Mase	/	mes	3	3.767		1	2	u Beste	3.767		1322	: 1	, 2		3.767	)	,	! 2 %	3	3.767	
1905	.47'	10	170	32	0	.08 0	.21	,59.	.45.	(E)	1901		1	!	(12)	C		,	ŀ	+ (N)	(38°)
1909	.11	,20 ,76 '	.19	.07	1	100	101 .	,26	0.	a.	1916						-				
1924	,01	123	,14	119	ļ	.07.	,33,	.05	(95)		1920	157	1.39	1.55	103	i i		1.78	1.04.		
1931	.N.	110	,30	0	3 4	119.	,3/*	,17	33	المالية المالية	1931	1.17	3.30.	1.37	148.	-	,0,0		i zari. 	1.5"-	
-3H3H	.57	87 .14	1112	70	42914	1.03	97	116	10 .18	en	E=11		1 5.25 Ohi	4.51	224	Was 11.	, 25	. 15°1	102 - 3H	202	146
1901	,03	Penney	fram in	1	(He)	.06	132.	Jaresen	02	400	1901	286.	167	343	130.	1400)					>
1416	122	143	1.40 .	11/5	(E)	.07	.11	9 :	101.	(J)	1916	131:	131	122 + 5 cm 4 m s	1.65	(g)	Xa	ret C	erole	na36	6
1924	1.10 .	.60	1.43 · ·81 ·67	183.	-	,09	70	00	.03	+	1920	1,71	1.83	3.14.	1.49	,	, 04	1:18	1.16	.04 .21 1.16	9
1937	114.	109.	4°7	19.	4	68	76	10	59	4	1931 j 31	1.42	223	2 68	3.14.	÷	1,26	2.79	1,7?	1,41	
	3,76	152	3.99	300	1500	1 085	-7.7:	92	.0.8		-314	7.48	Eat .		8.19 1.38	k., .4.1 - 1	142	193	,59		
1905	.99	1.97 ·	tand o	(4)4	390)		 Vurg		1	37050)	1901	, ,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	m	noun				( )	sense	7	(0)
1914	136	1.34	1.50	9		1071	D +	1.73	100	(A)	1909		512	"س.	1.37	390)	175	.47	.32 .48		(44°)
1924	178	120	1.73	(18) 1:11. 20		17/	1.44	1.32	1,00	. ()	*1 -	180.	1.84	3 64.	812'	(Z)	- 1	1.27	1,23	49 (-) 1	(A)
1935	1.31.	.3 /	. 32	.56		144	1.65	3.14:	2.37		د. الادام	2.14.	1.52	33.	1.20.		-	1.42.	,97 4,44.	238 . 1	
-14	2.86	6-21	2.12	134	4.00	4.87	3 97	6.93	285	1	حَدِّ	114	336	82	1.28	152	1.33		3 ( 2 )	1.38 %	
71501			,, .				.,,		1	1,2	Per 1901	ation of	og pour	<b>J</b> yr m		.995	17	2.79	3.67	.17	32-19
1905	1.150	1.30	thero	lina 33	3	Ta	tas:				1905	1.890	1.823 2.163 2.086	2.010	3.112	(F) In	diana	.84	1.39	(H)	
1916	1,92.	267 A	.60 N=0 A	14.27			, ~ 0	500	u	<i>!</i>	1916	1:233	2.183	2.064.	2096	<i>b</i>	32	.65	.84	60 .46 1.36	
1924 1947 1931	109	1.67	₹8 5.60.	.18		30	د ک	,12	12		1924	3 9829	1.1730	1.465	1.9530		202	2.11	10.8	1.25	
1935	6.59	797	. 83	3.95			1.2 9	2.67	.5 6	٠ ٧	435	1.850	1.998	9.706	10.765	础	93	1.45	.98	1.81	·
1901	(-1-5	133	1.26	66	6		·hs	Conguesty	30	(13)		1.952	2.04	1,941	2143		.04		77	70	
1904	. 8 5	.39	.76	369	(C)	, 34	.96.	1.70	* ,88	9.		Ken	tucky	37	030'		Colira	do ;	390	(N)	)
1920	1.20	120.	12.2	1.25.		. 36	.21 .	1.61 88	2.93		19 24		1		50 (	9)		,57	1.07	250	
1927	.60	1.05	2-/2	159		2.09.	.84	2.13.	1,29	!	1927	,5-1	1-04	131	1.07		.79	14.5	1.28	187	
-tit	3.15	265	450	343 -67	- (s)	00	.5-6	.96	39		1935	187	1,43	1.26	1,77	-	2.38	2.38	3.05	4.26	
1901		., .,				167	2.29	399	3.35					, ,	Ť		.,,				
1916	丰	landa	- 28	0 (	D	4		~ 3	30 T	)											
1924	1.35.	.82·		1 0.97		170	1/2.	.66	i									i	1	1	
1935	122	NO .	12 - 0	1.20		-1E 94	3.52	1,23	1.58.												
we	4,65	269	72	404		46	44	98	1.38												

					C	tres	ma	o Od	ist	20	- /	Bus	6	er I	whi					
	,	2	3	4	4,222		2	3	y	4,232	1	2	3		4.222					
		132	: 18.	.32	M)	, 45.	· 13	12/1	420	(Y)	0	2.86	1.83	3.40	1,20		Lentu	thy (3	703	
	120 120 1	119	100 007 117	.47.	123	123	.01.	109		-> 33	1.30	6020	1.83	131 124 1.55 4.72 1.41	1,83		.10	60	285	,11
2	128	.N	.15	.07		127	17 -13	113	.14		324	183	183/	2.68		1.07	.25	1.44	13/4	
	1.28	1,45	.98		elis dila	1.67	1.33	1.02	139	1 (I).	3.24	11.64	N. 89	17.07	(1) 467	1.07	2.55	2.03	1.13	
	. 1/3	1.80	1000	41', 094 1.10		.11	131	103	.07		147 3.79 2.12	1.01	196	181		1	anth C	eralin.	a (36)	
6	92 - -8/ 183 -98	1.43 .74 .62 1.49 .	154	1.18.	- 160	.03	10 L 13 123	.20	.06	-// 2	244 197 197	1.92 2.38	256	2.11 .48 .66 .85	,57	.92 .21 1.16	.87	1.18	1.16	
Pople	.19-	4:11 0 18 9 Maryl	202	173	of in all	.69	1.03	.39	41	aris at	1.81	11.94	8.18	8.83		2.29	1.26	2.79	1.77	
	1.34 .	0	6	.36	(M)		1.73	7,00	1.01		188	16	2.37		M		elar	do	390	
1 1	1.73 1 1.73 1	1.50	1.50	26 151 30 2901	O	1.32	12 /44	1.44	3211		2.37	3.64	1.37	134	1.84	.57	1.07	129	79	17
8	170 912 1.16	51RR 1.18	2,21	7,29	of ind	E: 5.11	389	296	4.89	1	1.20 1.35 3.52 1.76	1.45	289 1.45 44			2.14	2.66	4.31	2.07	
90	i	mes	i	369				H :			97	171 123	00 048	2:41 51	.42-					
9371		-60	1.05	2./2		2.15	· 88 ·/5 /-23	1.71	18	1.39	1.27	.53	0 0 1.08	0 0 0 9 7	,41					
30	.39 68 .92	.40	,50	30		1.03	1.17	3.30	1.59		238	.65	1.42	4.44						
	168	23.	150	.50		7.44	3.88	1.20	2.88		5.82	,27	4113 .46	8.47	D					
4 37	Ge.	ugia	33°	(	b)	Sa	with	Caroli	ma 3	3030	15.1	84	1.39	0						
26	·35	1.03	.38	·08 ·32 //23		2 88	1.79	1.67	3.60		.99	1.21	1.34		2.17					
38	1.78 2 15 272	1.43	3,52	1.63		.50	1.54	1.67	.83		254	4011	1,91	3.73						
100	. ,	Modua	der 4 3		m	取	endo	280		E	0	123	0	0	2)					
2/ 2/	.90	1.61	2930	36	184		1.97	.72	. 46		21	122	,24	138	).08					
30	1.29.	1.11	.44	2.13		1,20	1.07	1.87	192		23	0	0	2.78						
7 !	-	277	30	145	0	12	.53	162	389	(K) 8	.03	12,21	1:11	7.37						
30	.17	30	,53	152		1.04	.45		128	1,76										
26	7.13	87		3.40		5.5	1	t					!		-					

							1	TH	13.	LE	7	7		~ <i>/</i>	D	_					
				,	4	: 41	8-1	plan	- De	rior		tifle				<u></u>		1			
Bess	1	year 2	s agte	4	4.418	hris	300	is afi	us un b		P / /	3 <i>ers</i>	3	4	4,418		<u> </u>				
1899 1903 1907 1914 1916 1921 1925	132 120 1/1 0 109 109	1.27 1.27 1.37 1.36 1.14 1.18	.47 .14 .16 .19	.07	~14 ~14 ~.47	.59 .10 .09 .33	New .09 .10 .05 .07 .27 .19	90th 1/3 1/25 1/25 1/25 1/25 1/25 1/25 1/25 1/25	42 02 121 00 0 F 13 14	1	1897 13.7 1414 1425 1430	, 52	uda	286	1 / 39	(V)	.06	1 / K	53 201	.73	
1930 1934 1938 2 ave	.107	2.74	2:322	110		33	1.82	1 707 1 1194 1 1194	77.	+	1934 1938 2 ave	120 191 3.48 840	213	3.12	2.90	1	1 3.17	2256-	   4 31	160	3012
1899 1903 1907 1912 1916 1925 1930 1930 1938 5	194 192 160 183 144 119 4114 591	73 .60 1.43 .81 .62 1.44	10 1.40 1.40 1.44 1.44 1.64 1.64 1.64 1.64	1.27 1.14 1.18 1.67 1.77 40	-1.10	.07	news	.31	.58	03	1890 1903 1907 1907 1907 1907 1907 1935 1936 1936 1936 1936	2-15	117 144 117 145	.88. .67 .23 330 .2 6,22 1.256	12 171 137 137		1.36	000000000000000000000000000000000000000	.8 4 .0 .2 2 .1 4 .66 .0 .23	0038089 0 3500 35500 35500	.89
1499 1993 1993 1991 1916 1921 1930 1930 1938	36 1.74 0 1.11 .70 .44 3.97	1.42	140 2.99 400	4.87	L	41 140 2 39 .57	72 132 149 190	33 /2 /65 /47	30'P		1899 1413 190'Y 1915 1935 1935 1938 2001	994 179 1,29 5,21 1,04	12/1.6/188 1.11/0 3.81 .76	2.93 2.09 	30) (10 2.7 2.13 2.13 96 6.39	3)	VIA 2714 9 38 62 7 7 9 7	1.23 65 5.72	17 132 153 102 166 430 430 430	23 1000 14 5.89 7.740	48
7841 1903 1907 1912 1916 1924 1939 1938 8 8	.40 183 181 183 183 183 183 183 184 2017	1.10 1.34 1.44 1.83 3.24 2.42 7.01	1.30 1.22 1.50 1.22 1.49 1.91 2.23 1.87 1.87 1.87 1.87 1.87	1.41 .60 /83 .81 4.72 2.47 2.48 2.23	141	1.99 1.21 1.81. 244 .41 1.91 1.93 1.81 1.59	4.67 3.79 2.12 1.74 2.38 .60 1.196	1.57 1.67 1.07 1.07 1.07 1.07 1.060	179 196 196 196 196 196 196 196 196 196 19	1, 179 1,18 1,48	1930 1934 1938 2 2 1931 1935 1938 1938 2 2	174 91 46 557 557 2.08	1.47	1.20 1.20 1.20 1.20 1.20 1.20 1.23	52 2.16 2.67 179 N2 2.31	1.28	336 No 12/6 1.16 1.10 2.47 82	131 131 131 192 187 134 2114	.04	424 424 4.36 1:18 1:32 1:29 1:79 1:60	1.16
1403 1407 1912 1916 1921 1925 1930 1434 1938 E ave	1 2 2 2 17 1 2 4 1 2 0 1 3 3 - 1 1 1 8 1 2 9 7	3 4 × 3 4 × 3 × 4 × 4 × 4 × 4 × 4 × 4 ×	8-32 -90 /WZ 11-29 2.823			,83 ,20 ,39 ,39 ,68 -92 3 L8	0		1.05 1.05 1.30 1.71 1.678	- 200	Wisco Michi Iowa Massa	TATE	141 44 45 42 42	DE 3.70 Symior A Jn C Jn	37-YEAR  ool and Ideal H aly 26, aly 8, 1	OYGLE 4 Time 8 igh 1952 1 953 1 952 1	LES, FIG 4.222-YEA Symbol ar of Ideal * Dec. 26 F Oct. 4 I June 1 I June 1 I June 1 I Jan. 1	R CYCLE nd Time High 3, 1950 , 1951 5, 1951	4.418- Symbol of Id  A July B April C April D Jan.	1 7, 195 1 7, 195 17, 195	2 2 4
1899 1903 1907 1916 1921 1921 1934 1938	100 1/8 .50	268 154 .19 4.61	1.79 6 46 ,33	167 E 3 P.50	(3)	1.00 .39 .12 .30 .122 .152 .717	4.00 1.38 .62 .76	1.03 1.03 1.543	1.78 .72 .32 .423 1.93	-,e,z	Penns Ohio India Illin New Jo Maryl Color Eansa Kentu Virgi North Tenne Bouth Georg Texas Flori	oticut ylvania na ois ersey uri and ado s oky nia Carolin see Carolin	36 a 33 30 33 30 28	TRACE TO THE PROPERTY OF THE P	ug. 20, 0x 22, 1 3x, 20, 21, 21, 21, 21, 21, 21, 21, 21, 21, 21	953   1954   1952   1953   1955   195	Nov. 20   Nov. 20   Jan 28   1 Oot. 20   1 Feb. 1	1, 1964 7, 1951 1, 1953 4, 1952 1, 1952 0, 1951 1, 1952 3, 1951 25, 1951 1, 1950 6, 1951 1, 1953 14, 1962 1, 1952	E July F Aug. G Dec. H May I Dec. J June K Maro L June M Feb. M June O Marc O Marc P May Q Nov R Aug. S Aug. U Sept V Oct.	19, 195	1
										1	Vermo	nt ylvania	44 42 36	t to gro	wth and pril 3, st. 10, an. 14,	1952 ** 1951 ** 1953 **	r year re	sconcili	ation)	. 11, 19 16, 195 7 15, 195	


TABLE 11. EPOCHS OF 4.222-YEAR CYCLE: DOWNY AND HAIRY WOODPECKERS.

STATE	LATITUDE	SYMBOL	DOWNY WOODPECKER H	AIRY WOODPECKER
	0 1		Current Time	Current Time
			of Ideal High	of Ideal High
Alberta	54	A	Oct. 16, 1950	Nov. 2, 1952
British Columbia Washington	54	B W	Dec. 30, 1949 Sept 7, 1951	April 8, 1952
North Dakota	47 30 47 30	N N	Aug. 31, 1952	Nov 27, 1952 May 14, 1951
Quebec	47	Q ·	Dec. 10, 1951	April 15, 1952
Minnesota	47	M	April 25, 1951	Feb. 19, 1952
Montana	47	M	June 27, 1951	March 1, 1953
Maine South Dakota	44 30 44 30	M S	March 2, 1951	Nov. 20, 1953
New Hampshire	44	N	Oct. 7, 1951 Nov. 12, 1953	March 29, 1951 June 19, 1954
Vermont	44	V	April 8, 1952	Nov. 16, 1952
Ontario	44	0	April 8, 1952 Dec. 6, 1951	Dec. 1, 1951 Dec. 30, 1952
Wisconsin Oregon	44 44	W -	Dec. 8, 1953	Dec. 30, 1952
Michigan	43	M	Dec. 23, 1951 Oct. 15, 1952	Nov. 27, 1951 June 16, 1953
Iowa	42	Ï	Jan. 10, 1953	Oct. 29, 1953
Massachusetts	42	M	April 11, 1953	Sept. 27, 1950
New York Nebraska	42	N	Nov. 23, 1951	Sept. 3, 1950
Rhode Island	42 41 30	N R	Aug. 16, 1951 Nov. 20, 1951	Sept. 29, 1951 Nov. 12, 1950
Connecticut	41 30	C	Jan. 25, 1954	Sept. 27, 1950
Pennsylvania		P '	Oct. 18, 1951	July 7, 1950
	40	И	Jan. 3, 1954	April 11, 1950
Ohio Illinois	40 40	0 I	Nov. 16, 1951	Feb. 1, 1953
Indiana	40	Ī	July 30, 1952 Sept. 15, 1951	Sept. 8, 1952 Oct. 6, 1953
Maryland	39	M	Nov. 24, 1953	March 27, 1951
Missouri	39	M	Nov. 24, 1953 March 26, 1951	Nov. 21, 1951
Colorado	39	C	May 9, 1952	March 15, 1953
Kansas California	38 37 30	K C	Nov. 20, 1953 Dec. 27, 1951	March 2, 1951 July 26, 1951
Virginia	37 30	v	August 7, 1955	August 21, 1951
Kentucky	37 30	K	Jan. 7, 1955	August 25, 1950
North Carolina	36	N	Nov. 3, 1950	Sept. 4, 1951
Tennessee Oklahoma	36 35 30	T O	June 24, 1955 Dec. 26, 1950	June 13, 1951 April 7, 1951
	33 30	s	March 14, 1955	Sept 4, 1950
Georgia	33	G-	Jan. 8, 1955	Nov. 12, 1951
Mississippi	<b>3</b> 3	M	Nov 30, 1950	Tune 07 1054
Florida	28	F	June 27, 1955	June 27, 1954
TREE RINGS (subje	ect to gre	owth and	calendar year rec	onciliation)

TREE RINGS (subject to growth and calendar year reconciliation)

British Columbia	56	arrow	Oct. 15, 1952	Oct. 15, 1952
Vermont	44	7	*June 20, 1952	
Pennsylvania	. 42	8	August 26, 1952	*Oct. 5, 1954
Arizona	36	9	*May 10, 1954	*May 10, 1954
	(A star	* marks	a low)	

1															
	22 (	3	2.07	(R)	)			(	<u>ن</u> ((	کِ ۔	(	<u></u>		(	0) " 20
3	5	027	756:00	7.44	الإن ا	44, 44, 45, 45, 45, 45, 45, 45, 45, 45,	0 6 2	1.48	7 7 7 7	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	5.00	282	いないかい	2.49	1,2/ 1,3/2 1,3/2 1,0/2
hach	Μ.	Jan S	1030314	81:30		325	المرنية	10.	1/2,	£6 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5	4.87	المركبية	0000 V V V	2 2 6 7 7 5 7 5 6 7	1.000 0.000 0.000 0.000
90	7	Jedre :	1. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2.	\$ 15. C	. 7.	rous	20'4	20.4	立立	C 2 C 2 C 2 C 2 C 2 C 2 C 2 C 2 C 2 C 2	5.33	3000	2000	, ,	20000
25	` ~		7 12000	11.00 11.00	.32	25.5	54 C	रतेते.	13.7 Low	45.00522	4.4.4	108.	20.00 20.00	5 8 2º	3 0000
mour	33	.76	ά	(	0			(	( <u>z</u> )	09.		2	867	(	= = =
3	7 %	2/,	30 × 320 %	4.60	, é	24.	000	1.54	130	Sérisité à	2.92	€.	22. 48.	164	201-50
25	m . ş	00./	24.07 2007	5,2%	700	<b>9</b>	10,	30.	3	× 25.05.20	3.54	4539	2000	4.40	5 2 2 3 3 5
0 116	. N &	5.5.	222200	3.76	regar	ڊ د د	05°	.38	duga	S. S	× × × × × × × × × × × × × × × × × × ×	250	3200		و می در در
The state of the s	\ Z	بې ر	1 4 4 6 6 6 4 7 7 7 7 7 7 7 7 7 7 7 7 7 7	4.62	8	64.	8	يَرَبُ جُ	M	25.54.8 00.29.00	7,4,7,	7 6.	30.00	-2H 3.98	A-CC.
40 A	33	(2			0	0			(0)	7		3		. (3	33
4 6 4	<i>&gt;</i> ,	0,0	, 2, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0,	100	)\$ 0 3	1/0/0	10,0	1.86	4430	30405	15.5	440	123	28.	55.583
33	~	7	52 200	176	000	200	(S)	300	Robert	o varis s	2.23	yohio	insolvi.	12.5 2.6 2.6	ر در باهن
	1 %	nten	, ,,,, ,	136	<u>}                                    </u>	, w 0 /		9%;	The same	ないないか0	26%	E Jen	र्दहरी	1.68	66.7
		Z.	33.3	10 1 N	401	, av 5	10 50	1,44	75	3 2000	17 2.16 17.	The said	60/62 60/62	66	2023
	7/22	A)	.40		Q .	· ·			B			30,	بغ	et .	(B)
1	, y		4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4	2.72	history	.07	, ,	222	470301	i 40,2	00 CS	47030	204 5	1.05	470(
	2	5,40	36	1.04	Colum		05.	.34	کے	0000	% Q	3th	603	74.	١. ٤
	3	Ubenta	N. 4. 0 %	201	Buter	4 3	0000	.53	are home			the	133	1.65	Dueloc
	<i>-</i> ,			•				47	2			Ta		39.	
	Base	20061	\$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$	Sw?	1900	560	1930	3113	1900	0 2 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	GM3	00000	00000	Sen 3	98000



															-
	233	30'(R)	, h	0 %	90		0 3	0	>» >		2.2	3			6
	2	4/10	000,000	30,07	2 4 7 4 7 6	54.	200	100	2000	1,23	بالمنازية	183	25.0	200	33
	~	and	000,000	1 402 J	2000000	2 00 5	1	الم م الم	4 4 4 4	1.20	87.85	25.03	3. 2. X	3,75	.36
		ed se	0000000	.07 .00.	242514	81.	of the same	52.00	80:	00/	, 20 m	24.00	10 00 00 W	3.00	133
		32	10000000	83 g	\$ 0 mg = 0	10. 175	Den.	20.00	2 2 2 2	21:19	10 mg	22.	434	3.50	.43
9	4,222	B :	7.	4 =	4		•	7	7	(W	) 6 .	9		Book	.03
30 Buch	5	443	5.25.5.2	20 is	- 4 2 3 3 4 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5	18.	20 (1	. 5.5.	25.	222	4000	. 20	280	100 ×	200
-11	2	240	Susania de la seconda de la se	20 3	こうないまう	777	1 2 E	30%		2.29	2001	601	59.	Hart.	000
Having .	7	10 con	÷ 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	0	4 3 3 3 4 4	27.2	斗	, 4 6	. 67.	2.39 ,60	2040	20.	70°	New	× 00 %
10 m		A 60.	こうかなからだか	22 8	0 2,20	180		34.	2000	43.72	2000	200	000	.07	50.
and	4,222				٤,		9	04	- 1	3	)	and a surface has abordered a			, h.,
131 E /	7	(X)	400	si is	0000	20	44030	· 8 %.	1.76.	85.	24.	0,00	3.	خ 🗷	7.0
4 131 year Park	7	25		12 27	5000	7	bota	727	\$ 2.4	125	なったい	0000	87.	440	00
The state	~	Mortan	12%	17 137 Mary	10%	50.	South De	5,7.5	85.00	\$ £' \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$	000	0000	.00.		374
13	1	M	4:4	31. 3	\$ 00	00.	Sew	72	1.330	141.98	0 80	0000	42.	Very	0
	4222				64.			07					77	1	
	5 >	: (2)	, dans	5,08	@ \$ 50°	9 35		(10		2500		1	80.		.94.
	Sm	4703	100	.02 .08	100%	5 0%	3 ú	3	18,	20%	470 (4)	9/.	2000	0	.26.
	No or	hungton	30000		3,3		7	24	67	7.25	whee c	7.00	40. 27.	water	· 70.
		Te .	( ) 6 9 6 W	Bretsh :	3.2	० र छ व यक र ०	3	3	37.00	2 2 2	3	, ,	9040	3	.67
	Sea Sea	h061	00000000 0000000 000000000000000000000	1900 1900 1900	1927	1935 1935 1935	1900	6969	388		19000	222	1938	1900	1909

(3							**********												
			7.		(0)	) 10.			(3)		2								
4	302	1.00/	250,000	000 2	030'	ō	1000	14.	30,	7 7	,	1,20							
	5.5	676	2 2 6 5 2 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5	00 7	a 37		20.	00 0 00 0	376	מלים	0 100	0; Qu							
	4.24	and	00000	2000		200	1000	000	in	. c. j.	100 or 600	50.	American (1)						
.23	4.71	Indan	30 804	30 X	5 3	9	1000	23.	UNA	100	127	2000							
er ce		6	<i>711.</i>			1/3	)				<i>k</i> ,							, ,	
	25-	120		247.	200		0.38	90	0	0 0		17.	1386	327	9	149	5) 67:	17, 17, 17, 19, 19, 19, 19, 19, 19, 19, 19, 19, 19	0-1
	1 60:	-		20%	.,0	13.	200.	200.	ohh	00	, , ,		Smer	500		1.62	-	50 00 50 00	70
. 0.16	50.	Nebrasha		225 2	*	00 0		200			0 6	0 10	Conol	نا ما			1 62	977	
	a. 1	×.		,	1 de		500		and	03 60			\$ -			200	7. 78	11.	7.5%
				it it	X	.,,	THE STATE ST										-	.43	
iner -		75.	ų,		8	. 2 .			6	~		201	P			0,0			29
	97.	(a)	4:050	10%	370	1.14	.63	1,64	360		325	35.	360	04. CC:	77	35.030	.26	1.05 5.5.5 4.8	2,82
1 W	20.	940	40 24 7	900	oper 1	778	12.2	3.	rolina	00	0,230	2000	are	0 44	0 00	0 2 5	15	30 30 30 30 30 30 30 30 30 30 30 30 30 3	200
100	zi.	, 60°	\$ 65.50 P	\$00 ox/	3	, o, n,	0,2	13.	roch	76'	01.00		wer	2000	100	10%	71.	1 1 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	224
. 36	11. 12	Pa B	zoża:		le.	33.	ייי ייי	## . 200 . 7 . 7 . 7 . 7 . 7 . 7 . 7 . 7 . 7 .	8	15	800	3000	19	22.	,20 .13	1.40 2.50 Blog	.30	1. 25 S	357
		B		- ,	1			7"				31 (		0.00		3	Ç	43	
	29.	8	33.05	1.80	n e	1000	1001	, in	E) % %	39.	27.	2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	(0)	0.000	0,0		10.	707	300
-	17.3	the state of	4,5,5			4	.13.				46.		-	1 1		W 260	رغ	نه نه نه نه م م م م ی پر یام م	شرشر
-	109	Dal	32.75		139 mlan 23		24.		, 3	46.00	0000	3. 402	3	9/:	0000	25. 4	7:	マッシッション マッシッシッシッシッシッション	3.34
47	- 37. 1.69 	Nanth	2326	2 2 3	多艺	2000			3	. 520		77.77		-/-	10 m	3	3.4.	\$ 45 m	35.7
-	17 F			7 77			2000	O. F			190:	* # # C	03	WW. 25.78	22.28		5702	-1	
26	Sw Y	900	2000	322	300	1904	8 420	8 8	700	263	1921	Se my	900	1917	6,6,00	w \$ 0	2002	1926	10 m 3

